

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Amendments to the Claims

1. through 7. (Cancelled)


8. (Withdrawn) Di(1-1-isopropyl-3-methylbut-2-enyl)borane of the formula (Ia).

9. (Withdrawn) A bis(allyl)borane of the formula (I) obtainable by a process as claimed in claim 1.

10. (Withdrawn) A Suzuki coupling reaction product obtained through use of a bis(allyl)borane of the formula (III) or (V) in C-C coupling reactions





11. (Currently Amended) A process for preparing ~~boronic acids by reaction of a bisallyl alkylboronate comprising the steps of reacting a diene with sodium borohydride in the presence of an oxidant to form the corresponding bis(allyl)borane of the formula (I) as described in claim 1~~





where R^1 - R^6 are H, aryl or substituted or unsubstituted C_1 - C_4 -alkyl or two radicals R may be closed to form a cyclic system.

and further ~~reaction of reacting~~ the borane (I) with an appropriate alkene (II) or alkyne (IV) to give the



alkylbis(allyl)borane (III) or ~~alkenylbis(allyl)borane (V)~~



wherein R⁹ to R¹² are selected from the group consisting of aryl, substituted or unsubstituted, alkyl-(C₁-C₈), branched and/or substituted alkyl-(C₁-C₈), alkoxy-(C₁-C₈), acyloxy-(C₁-C₈), Ophenyl, fluorine, chlorine, NO₂, NH₂, NHalkyl-(C₁-C₈), Nalkyl-(C₁-C₈), CN, CHO, SO₃H, SO₃R, SO₂NH₂, SO₂N(alkyl-(C₁-C₈))₂, SO₂-alkyl-(C₁-C₈), COO-alkyl-(C₁-C₈), CONH₂, CO-alkyl-(C₁-C₈), NHCHO, CF₃, 5-membered heteroaryl and 6-membered heteroaryl, where two radicals may also form a cyclic ring system which may contain heteroatoms.

~~which is oxidized and oxidizing~~ directly in the presence of an oxidant to form the corresponding bisallyl alkylboronate or alkenylboronate and, if desired, subsequent conversion into a derivative.

12. (Cancelled)

13. (Original) The process as claimed in claim 11, wherein the oxidant used is formaldehyde, acetone, glyoxal or diacetyl.

14. (Withdrawn) A Suzuki coupling reaction product obtained by using bis(allyl) alkylboronate or alkenylboronate produced as claimed in claim 11 in C-C coupling reactions.